



ELSEVIER



<https://doi.org/10.1016/j.ultrasmedbio.2020.04.033>

● *Clinical Note*

## LUNG ULTRASOUND IN PATIENTS WITH ACUTE RESPIRATORY FAILURE REDUCES CONVENTIONAL IMAGING AND HEALTH CARE PROVIDER EXPOSURE TO COVID-19

SILVIA MONGODI,<sup>\*</sup> ANITA ORLANDO,<sup>\*</sup> ERIC ARISI,<sup>\*</sup> GUIDO TAVAZZI,<sup>\*,†</sup> ERMINIO SANTANGELO,<sup>‡</sup>  
 LUCA CANEVA,<sup>\*</sup> MARCO POZZI,<sup>\*</sup> ELEONORA PARIANI,<sup>†</sup> GIADA BETTINI,<sup>†</sup> GIUSEPPE MAGGIO,<sup>\*</sup>  
 STEFANO PERLINI,<sup>‡,§</sup> LORENZO PREDÀ,<sup>†,¶</sup> GIORGIO ANTONIO IOTTI,<sup>\*,†</sup> and FRANCESCO MOJOLI<sup>\*,†</sup>

<sup>\*</sup> Anaesthesia and Intensive Care, San Matteo Hospital, Pavia, Italy; <sup>†</sup> Department of Clinical-Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Pavia, Italy; <sup>‡</sup> Department of Translational Medicine, Università del Piemonte Orientale, Novara, Italy; <sup>§</sup> Emergency Medicine, San Matteo Hospital, Pavia, Italy; and <sup>¶</sup> Radiology Department, San Matteo Hospital, Pavia, Italy

(Received 18 April 2020; revised 28 April 2020; in final form 29 April 2020)

**Abstract**—Lung ultrasound gained a leading position in the last year as an imaging technique for the assessment and management of patients with acute respiratory failure. In coronavirus disease 2019 (COVID-19), its role may be of further importance because it is performed bedside and may limit chest X-ray and the need for transport to radiology for computed tomography (CT) scan. Since February 21, we progressively turned into a coronavirus-dedicated intensive care unit and applied an ultrasound-based approach to avoid traditional imaging and limit contamination as much as possible. We performed a complete daily examination with lung ultrasound score computation and systematic search of complications (pneumothorax, ventilator-associated pneumonia); on-duty physicians were free to perform CT or chest X-ray when deemed indicated. We compared conventional imaging exams performed in the first 4 wk of the COVID-19 epidemic with those in the same time frame in 2019: there were 84 patients in 2020 and 112 in 2019; 64 and 22 (76.2% vs. 19.6%,  $p < 0.001$ ) had acute respiratory failure, respectively, of which 55 (85.9%) were COVID-19 in 2020. When COVID-19 patients in 2020 were compared with acute respiratory failure patients in 2019, the median number of chest X-rays was 1.0 (1.0–2.0) versus 3.0 (1.0–4.0) ( $p = 0.0098$ ); 2 patients (3.6%) versus 7 patients (31.8%) had undergone at least one thoracic CT scan ( $p = 0.001$ ). A self-imposed ultrasound-based approach reduces the number of chest X-rays and thoracic CT scans in COVID-19 patients compared with patients with standard acute respiratory failure, thus reducing the number of health care providers exposed to possible contamination and sparing personal protective equipment. (E-mail: [silvia.mongodi@libero.it](mailto:silvia.mongodi@libero.it)) © 2020 World Federation for Ultrasound in Medicine & Biology. All rights reserved.

**Key Words:** Lung ultrasound, Lung monitoring, COVID-19, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), ARDS.

### INTRODUCTION

On March 11, 2020, the World Health Organization declared the new coronavirus disease 2019 (COVID-19) as an ongoing pandemic emergency. The first reported case had been identified in Wuhan, China, on December 8, 2019. Since then, the disease has rapidly spread worldwide. Around 5%–10% of COVID-19 patients require mechanical ventilation for acute respiratory failure (ARF), of whom 65%–85% are classified as having acute respiratory distress syndrome (Wu and McGoogan, 2020; Yang et al. 2020). The typical radiologic findings are consolidations and ground glass opacities with a peripheral distribution. Frequently, lung involvement is bilateral, with destruction of pulmonary parenchyma

Address correspondence to Silvia Mongodi, Rianimazione I, Fondazione IRCCS Policlinico S. Matteo, DEA piano-I, Viale Golgi 19, 27100 Pavia. E-mail: [silvia.mongodi@libero.it](mailto:silvia.mongodi@libero.it)

On behalf of the Group for Research in Intensive Care in Pavia (the GRIP): Francesco Mojoli, Silvia Mongodi, Anita Orlando, Marco Pozzi, Guido Tavazzi, Eric Arisi, Luca Caneva, Giuseppe Maggio, Luca Civardi, Giuseppe Sala Gallini, Roberta Puce, Federico Visconti, Fiorenza Ferrari, Alessandro Amatu, Andrea Stella, Andrea Colombo, Silvia Bonaiti, Giulia Salve, Sandro Pregnotato, Margherita Cavagnino, Eleonora Pariani, Francesco Daverio, Giada Bettini, Gilda Maria Siano, Cinzia Villa, Valentino Dammasa, Erminio Santangelo.

On behalf of the Pavia COVID-19 Task Force: Carlo Marena, Monica Calvi, Giuseppina Grugnetti, Marco Maurelli, Alba Muzzi, Raffaele Bruno, Paolo Lago, Gianluigi Marsaglia, Stefano Perlini, Alessandra Palo, Fausto Baldanti, Luigi Oltrona Visconti, Angelo Guido Corsico, Antonio Di Sabatino, Francesco Mojoli, Giorgio Iotti, Marco Benazzo, Carlo Nicora, Antonio Triarico, Vincenzo Petronella.

caused by consolidations and interstitial inflammation (Chung *et al.* 2020).

As in all acutely ill respiratory patients, lung imaging is mandatory for assessment of disease severity and to guide clinical management. However, radiology societies have advised against the systematic use of computed tomography (CT) and have suggested reducing the number of chest X-rays (CXRs) in COVID-19, to minimize the exposure of health care providers to this highly infective disease (Simpson *et al.* 2020). In this context, lung ultrasound (LUS) could be the ideal imaging technique as it is available at the bedside, it is performed directly by the in-charge physicians and it provides reliable information for diagnosis and monitoring of ARF (Chiumello *et al.* 2018; Mojoli *et al.* 2019; Pescarissi *et al.* 2020). The aim of our study was to test whether the systematic use of a LUS-based approach could reduce the number of conventional radiologic exams during the COVID-19 epidemic.

## METHODS

We admitted the first COVID-19 patient on February 21 and progressively transformed our intensive care unit (ICU) into a coronavirus unit, reaching 100% of patients with COVID-19 on March 6, 2020 (Mojoli *et al.* 2020). Because our health care providers have high skills and experience in LUS and habitually use ultrasound in daily management of ARF patients, although in general integrated with traditional imaging, from the beginning we decided to use an LUS-based approach. The aim of this self-imposed limitation was to minimize transport of COVID-19 patients to the radiology department, reduce the number of bedside CXRs and therefore reduce the exposure of health care professionals. A complete 12-area examination, with computation of the LUS score (Mongodi *et al.* 2017), is performed daily with a Vivid iq ultrasound machine (GE Healthcare, Chicago, IL, USA). A 9-MHz linear probe is used if the pleural line is visualized with one focus on the pleural line, artifact-erasing software and harmonics are abolished, and depth is adjusted to be at least twice the depth of the pleural line (Mongodi *et al.* 2019); a phased-array probe is used in case of consolidations or pleural effusions. The complete LUS examination is finalized to monitor lung aeration; guide the respiratory strategy, that is, positive end-expiratory pressure titration, pronation (Bouhemad *et al.* 2011; Mojoli *et al.* 2019); and identify complications (*i.e.*, pneumothorax and ventilator-associated pneumonia [Mongodi *et al.* 2016; Mojoli *et al.* 2019]). On-duty physicians are free to perform CT or CXR when deemed necessary. Operators were either recognized experts in the field or trainees who had completed 25 supervised examinations (Rouby *et al.* 2018).

We computed the number of traditional imaging exams performed in all ICU patients, including those previously admitted, from February 22 to March 22, 2020, and compared it with the number for the same time frame in 2019. We focused on CXRs and chest CTs per patient and per patient-bed days, in particular for patients with ARF. Informed consent was collected according to the *ad hoc* procedures defined by the ethics committee for the COVID-19 pandemic. The treating physicians had the responsibility for the patients' data management and protection aiming at the improvement in treatment and safety.

Quantitative and categorical variables are expressed as the median and number (percentage), respectively. Normally distributed data were assessed with the Shapiro–Wilk test. Comparisons between time frames were performed with the unpaired Wilcoxon–Mann Whitney *U*-test for quantitative variables and the Fisher exact/ $\chi^2$ -test for categorical variables. Comparisons were performed between the following pairs: overall ICU population in 2020 versus 2019; ARF patients in 2020 versus those in 2019; COVID-19 patients in 2020 versus ARF patients in 2019. A *p* value <0.05 was considered to indicate significance. Statistical analysis was performed using STATA14 for Macintosh.

## RESULTS

Over the time frames analyzed, we had 84 patients in 2020 and 112 in 2019 in our 23-bed ICU, for a total of 969 and 1070 and patient-bed days, respectively (Table 1); we had an increased number of ARF patients in 2020 (76.2% vs. 19.6%, *p* < 0.001), mainly because of COVID-19 (85.9%). Comparison of ARF patients revealed that male sex was more frequent (53 [82.8%] vs. 12 [54.6%], *p* = 0.008) and body mass index was higher (27.5 [24.6–30.9] vs. 23.5 [20.1–27.6], *p* = 0.0132) in 2020 than 2019.

Results concerning the use of traditional imaging are outlined in Table 2. No significant difference was observed when comparing the overall ICU population in 2020 versus that in 2019. When comparing COVID-19 ARF in 2020 with ARF in 2019, the percentages of patients having undergone at least one CXR were similar (85.5% vs. 86.4%, *p* = 1.000), but the median number of CXRs per patient was significantly lower in COVID-19 (1.0 [1.0–2.0] vs. 3.0 [1.0–4.0], *p* = 0.0098). Similar results were obtained when considering the number of exams performed per patient-bed day. The percentage of patients who underwent at least one thoracic CT scan was lower in COVID-19 versus ARF in 2019 (3.6% vs. 31.8%, *p* = 0.001). Only 2 COVID-19 patients underwent CT, one twice. Similar results are obtained when considering the median number of CT scans per patient and per patient-bed day.

Table 1. Clinical features in the ICU populations during COVID-2019 in 2020 and during the same time frame in 2019\*

Characteristic	2019 (112 patients)	2020 (84 patients)	<i>p</i> Value
Males	63 (56.3)	65 (77.4)	<b>0.002</b>
Age, y	64.0 [53.0–74.0]	63.5 [50.5–70.0]	0.2726
BMI, kg/m <sup>2</sup>	25.8 [22.8–30.0]	26.3 [23.5–30.7]	0.1774
ICU stay, h	74.0 [23.0–239.0]	221.5 [111.5–341.5]	< <b>0.0001</b>
Provenience			
Emergency department	46 (41.1)	19 (22.6)	< <b>0.001</b>
Subacute care	10 (8.9)	32 (38.1)	< <b>0.001</b>
Surgical ward	39 (34.8)	9 (10.7)	< <b>0.001</b>
Medical ward	17 (15.2)	24 (28.6)	< <b>0.001</b>
Post-surgical patients	36 (32.1)	11 (13.1)	<b>0.002</b>
Acute respiratory failure	22 (19.6)	64 (76.2)	< <b>0.001</b>
COVID-19	0.0 (0.0)	55 (65.5)	
Males	12 (54.6)	53 (82.8)	<b>0.008</b>
Age, y	63.0 [50.0–73.0]	62.5 [51.5–70.5]	1.0000
BMI, kg/m <sup>2</sup>	23.5 [20.1–27.6]	27.5 [24.6–30.9]	<b>0.0132</b>
ICU stay, h	210.0 [105.0–522.0]	238.0 [151.0–378.5]	0.9882

COVID-19 = coronavirus disease 2019; ICU = intensive care unit; BMI = body mass index; IQR = interquartile range.

\* Values are expressed as the number (%) or median [interquartile range]. Significant *p* values < 0.05 are in boldface.

## DISCUSSION

The main result of the present work is that a self-imposed LUS-based approach reduces the number of CXRs and thoracic CT scans in COVID-19 patients compared with standard ARF patients; this implies that LUS helps to reduce the need for transport of infectious patients to the radiology department and for bedside CXRs, thus reducing the number of health care providers exposed to possible contamination and sparing personal protective equipment.

Interestingly, in our center, not all patients with ARF underwent at least one CXR and fewer than 1 in 3 underwent chest CT in the examined time frame in 2019. For many years, LUS has been used in our center for the

assessment and management of ARF patients. Even in the context where the use of conventional chest imaging had already been optimized by the daily practice of LUS, a self-imposed LUS-based approach could further reduce the number of conventional exams. The two COVID-19 patients who underwent CT for lung assessment both had a condition preventing LUS examination (*i.e.*, subcutaneous emphysema and pneumomediastinum).

As for limitations, first we analyzed only thoracic CT scans and CXRs, as no COVID-19 patient required imaging other than thoracic, so far. Second, the integration of LUS findings in the management of ARF requires well-trained physicians (Rouby et al. 2018). Automation may be of help in the future to overcome this limitation

Table 2. Use of traditional radiology in the ICU populations during COVID-19 epidemic in 2020 and in the same time frame in 2019\*

	Chest X-ray			Thoracic CT scan		
	2019	2020	<i>p</i>	2019	2020	<i>p</i>
Patients who underwent traditional imaging						
Overall	78 (69.6)	68 (81.0)	0.072	18 (16.1)	14 (16.7)	0.911
With ARF	19 (86.4)	53 (82.8)	1.000	7 (31.8)	8 (12.5)	<b>0.039</b>
COVID-19		47 (85.5)	1.000 <sup>†</sup>		2 (3.6)	<b>0.001<sup>†</sup></b>
No. of imaging exams per patient						
Overall	1.0 [0.0–3.0]	1.0 [1.0–2.0]	0.6672	0.0 [0.0–0.0]	0.0 [0.0–0.0]	0.8928
1.8 ± 2.1	1.6 ± 1.4			0.2 ± 0.5	0.2 ± 0.5	
With ARF	3.0 [1.0–4.0]	1.0 [1.0–2.0]	<b>0.0079</b>	0.0 [0.0–0.1]	0.0 [0.0–0.0]	<b>0.0372</b>
3.1 ± 2.7	1.5 ± 1.1			0.4 ± 0.6	0.2 ± 0.5	
COVID-19 <sup>†</sup>		1.0 [1.0–2.0]	<b>0.0098<sup>†</sup></b>		0.0 [0.0–0.0]	<b>0.0006<sup>†</sup></b>
1.5 ± 1.1					0.06 ± 0.3	
Exams per patient-bed day						
Overall	204 (19.1)	133 (13.7)	<b>0.001</b>	23 (2.2)	18 (1.9)	0.639
With ARF	69 (19.1)	98 (12.4)	<b>0.003</b>	11 (3.0)	11 (1.4)	0.058
COVID-19 <sup>†</sup>		85 (13.8)	<b>0.028<sup>†</sup></b>		3 (0.5)	<b>0.001<sup>†</sup></b>

ARF = acute respiratory failure; COVID-19 = coronavirus disease 2019; ICU = intensive care unit.

\* Values are expressed as the number (%), median [interquartile range] or mean ± standard deviation. Significant *p* values < 0.05 are in boldface.

† COVID-19 patients in 2020 are compared with ARF patients in 2019.

for lung aeration quantification (Brusasco *et al.* 2019; Piculjan *et al.* 2019).

## CONCLUSIONS

A LUS-based approach for the management of COVID-19 ARF reduces the number of conventional thoracic imaging exams and reduces exposure of health care professionals.

*Conflict of interest disclosure*—S.M. received fees for lectures from GE Healthcare, outside the present work. A.O. received fees for manuscript preparation from Hamilton Medical, outside the present work. M.P. received fees for lectures from Hamilton Medical, outside the present work. L.C. received fees for lectures by GE Healthcare, outside the present work. G.T. received fees for lectures by GE Healthcare, outside the present work. G.A.I. received fees for lectures by Hamilton Medical, Eurosets, Getinge, Intersurgical SpA, and Burke & Burke, outside the present work. F.M. received fees for lectures from GE Healthcare, Hamilton Medical, and SEDA SpA, outside the present work. There is an active research agreement between University of Pavia and Hamilton Medical, outside the present work.

## REFERENCES

- Bouhemad B, Brisson H, Le-Guen M, Arbelot C, Lu Q, Rouby JJ. Bed-side ultrasound assessment of positive end-expiratory pressure—induced lung recruitment. *Am J Respir Crit Care Med* 2011;183:341–347.
- Brusasco C, Santori G, Bruzzo E, Trò R, Robba C, Tavazzi G, Guarracino F, Forfori F, Boccacci P, Corradi F. Quantitative lung ultrasonography: A putative new algorithm for automatic detection and quantification of B-lines. *Crit Care* 2019;28(23):288.
- Chiumello D, Mongodi S, Algieri I, Vergani GL, Orlando A, Via G, Crimella F, Cressoni M, Mojoli F. Assessment of lung aeration and recruitment by CT scan and ultrasound in acute respiratory distress syndrome patients. *Crit Care Med* 2018;46:1761–1768.
- Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, Cui J, Xu W, Yang Y, Fayad ZA, Jacobi A, Li K, Li S, Shan H. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 2020;295:202–207.
- Mojoli F, Bouhemad B, Mongodi S, Lichtenstein D. Lung ultrasound for critically ill patients. *Am J Respir Crit Care Med* 2019;199:701–714.
- Mojoli F, Mongodi S, Grugnetti G, Muzzi A, Baldanti F, Bruno R, Tricarico A, Antonio Iotti G. Setup of a dedicated coronavirus intensive care unit: Logistical aspects [e-pub ahead of print]. *Anesthesiology* 2020. doi: 10.1097/ALN.0000000000003325 Accessed Mar 31.
- Mongodi S, Via G, Girard M, Rouquette I, Misset B, Braschi A, Mojoli F, Bouhemad B. Lung ultrasound for early diagnosis of ventilator-associated pneumonia. *Chest* 2016;149:969–980.
- Mongodi S, Bouhemad B, Orlando A, Stella A, Tavazzi G, Via G, Iotti GA, Braschi A, Mojoli F. Modified lung ultrasound score for assessing and monitoring pulmonary aeration. *Ultraschall Med* 2017;38:530–537.
- Mongodi S, Stella A, Orlando A, Mojoli F. B-Lines visualization and lung aeration assessment: Mind the ultrasound machine setting. *Anesthesiology* 2019;130:444.
- Pescarissi C, Brogi E, Sidoti A, Corradi F, Forfori F. Role of lung ultrasound in the peri-operative evaluation of surgical patients. *Minerva Anestesiol* 2020. doi: 10.23736/S0375-9393.20.14417-1.
- Piculjan A, Sustic M, Brumini G, Kuaric J, Sustic A. Reliability of B-line quantification by different-level observers and a software algorithm using point-of-care lung ultrasound. *J Clin Monit Comput* 2019;. doi: 10.1007/s10877-019-00440-7.
- Rouby JJ, Arbelot C, Gao Y, Zhang M, Lv J, An Y, Wang C, Bin D, Barbas CSV, Dexheimer Neto FL, Prior Caltabeloti F, Lima E, Cebey A, Perbet S, Constantin JM, APECHO study Group. Training for lung ultrasound score measurement in critically ill patients [e-pub ahead of print]. *Am J Respir Crit Care Med* 2018. doi: 10.1164/rccm.201802-0227 LE, Accessed Mar 20.
- Simpson S, Kay FU, Abbara S, Bhalla S, Chung JH, Chung M, Henry TS, Kanne JP, Kligerman S, Ko JP, Litt H. Radiological Society of North America expert consensus statement on reporting chest CT findings related to COVID-19. Endorsed by the Society of Thoracic Radiology, the American College of Radiology, and RSNA [e-pub ahead of print]. *Radiol Cardiothorac J Thorac Imaging* 2020. doi: 10.1097/RTL.0000000000000524 Apr 21 Accessed.
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: Summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323:1239–1242.
- Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T, Wang Y, Pan S, Zou X, Yuan S, Shang Y. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: A single-centered, retrospective, observational study [e-pub ahead of print]. *Lancet Respir Med* 2020. doi: 10.1016/S2213-2600(20)30079-5 Accessed Feb 24.